## SACOX 60 - salinomycin powder

Huvepharama, Inc.

### FRONT PANEL

Brand of salinomycin sodium
Type A Medicated Article
For use in broiler, roaster, replacement
(breeder and layer) chickens, and quail only

#### ACTIVE DRUG INGREDIENT

Salinomycin equivalent to 60 grams salinomycin sodium activity per pound

For use in manufacturing medicated feeds only

#### OTHER INGREDIENTS

Calcium Carbonate, Mineral Oil, Calcium Silicate and Silicon Dioxide

## INDICATIONS FOR USE

# Broiler, roaster and replacement (breeder and layer) chickens

Use Sacox 60 to provide salinomycin sodium concentrations ranging from 40-60 g/ton (0.0044% to 0.0066%) for the prevention of coccidiosis in broiler, roaster and replacement chickens caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions.

Quail

Use Sacox 60 to provide salinomycin sodium concentrations of 50 g/ton (0.0055%) for the prevention of coccidiosis in quail caused by *Eimeria dispersa* and *E. lettyae*.

### MIXING AND FEEDING DIRECTIONS

Mixing and Feeding Directions

Mixing Directions

Thoroughly mix Sacox® 60 with non-medicated feed to provide the

level of salinomycin sodium per ton as indicated in the table below.

Salinomycin Sodium	Amount of Sacox® 60 Type A Medicated	
Activity	Article (60	0 g/lb) Per Ton of Feed
(grams per ton)	Lbs	Grams
40	0.67	304
45	0.75	341
50	0.83	377
55	0.92	418
60	1.00	454

Feed continuously as the sole ration

**Important** 

Must be thoroughly mixed in feeds before use. Not approved for use with pellet binders.

## **CAUTION**

CautionCaution

May be fatal if fed to adult turkeys or to horses

## WARNING

Warning

Do not feed to laying hens producing eggs for human consumption. When mixing and handling Sacox 60 premix, use protective clothing, impervious gloves, eye protection, and an approved dust respirator. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water

## STORAGE AND HANDLING

Storage and Handling

Store in a cool, dry place

ANADA #200-075, Approved by FDA

Distributed by:

Huvepharma, Inc.

Peachtree City, GA 30269

Net Wt. 50 lb. (22.68 kg)

# **BAG LABEL IMAGE**

Bag Label Image

